

Proton Chemical Shifts in NMR. Part 12¹. Steric, electric field and conformational effects in acyclic and cyclic ethers.

Raymond J. Abraham,*^a Mark A. Warne^a and Lee Griffiths^b

^a Chemistry Department, The University of Liverpool, P.O.Box 147, Liverpool L69 3BX

^b Zeneca Pharmaceuticals, Mereside, Alderley Park, Macclesfield, Cheshire, SK10 4TG

The proton resonance spectra of tetrahydropyran at room temperature and -85⁰ C where the ring inversion is slow on the NMR time scale, 2-methoxy, 3-methyl, and several 4-substituted tetrahydropyrans, 2-methyl-1,3-dioxolane and the rigid cyclic ethers 7-oxa-bicyclo-[2.2.1]-heptane and 1,8-cineole were recorded and completely analyzed.

These results together with literature data on acyclic and cyclic ethers (1,3- and 1,4-dioxane, dioxolane, 4-oxa-5 α -androstane etc.) allowed the determination of the oxygen substituent chemical shifts (SCS) in these systems. This data set consisting of 78 proton chemical shifts in seventeen compounds was then used to test the application of a previous theoretical model of proton chemical shifts to these compounds.

It is shown that the model gives a very good account of the proton chemical shifts in these systems. The ether oxygen SCS are due to both steric and electrostatic terms, the steric term predominating at short distances (e.g. in the 1,3-diaxial interactions in methoxy-cyclohexanes).

Conformational isomerism in these compounds was also investigated. Low temperature NMR gave ΔG (eq-ax) + 1.0 kcal mol⁻¹ for 4-hydroxy THP. Analysis of the couplings in the CH₂OH side chain of 2-hydroxymethyl THP showed that the preferred conformer was gt in both chloroform and acetone solvents.

Introduction

Ethers and alcohols are of very considerable practical and theoretical importance in chemistry and commerce. Yet although the effect of the electronegative oxygen atom on proton chemical shifts has been known for *ca* five decades² a definitive analysis of oxygen substituent chemical shifts (SCS) in proton spectra has not been performed to date despite a number of investigations^{3a-c,4}. Zurcher^{3a} was limited to observing only the methyl groups in steroids in his


pioneering studies but concluded that the C-O bond anisotropy was not important for the OH group SCS. Schneider et al^{3b} regarded the electric field term as the dominant term but Hall^{3c} suggested that the chemical shift difference between the anomeric protons of the C₂-O axial and C₂-O equatorial sugars could be accounted for by C-O anisotropy alone. Yang et al⁴ concluded that electric field, anisotropy and a constant term were necessary to reflect the observed ether SCS in oxasteroids but did not consider any steric contributions.

The major reason for the absence of any systematic investigation is simply the divalent nature of the oxygen atom which gives an additional degree of freedom when compared to monovalent substituents. For example the SCS of the hydroxyl group may well be dependent on the position of the hydroxyl proton. The CH.OH coupling in alcohols shows that the OH proton is usually not in a single orientation but may have a preferred conformation⁵. However the precise populations of the different conformers cannot be estimated accurately. For these reasons we have restricted this initial investigation of oxygen SCS to ethers. Even in ethers the conformational mobility caused by the oxygen atom produces considerable problems. The only simple acyclic ethers that exist in one conformation are dimethyl ether and methyl-*t*-butyl ether. Methyl-ethyl ether has two populated conformers, diethyl ether four and the number of conformations of more complex ethers (e.g. diglyme, 2-methoxy-ethyl ether) is prohibitive for any quantitative calculation.


Thus we have selected mainly cyclic ethers of known conformation. The proton chemical shifts in 1,3- and 1,4-dioxanes and their methyl derivatives have been obtained in a number of previous investigations⁶⁻⁹. In contrast the tetrahydropyran (THP) ring system has been mainly studied in investigations of the substituent ΔG° values by low temperature ¹³C NMR^{7,10}. The proton spectrum of THP (**1**) has been recorded at low temperature by a number of investigators¹¹⁻¹⁵, but the full assignment has not been given. Gatti et al.¹¹ decoupled H3/H4 and observed the H2 AB pattern at 100 MHz. Lambert et al.¹² examined deuterated analogues and reported the ΔG^\ddagger and coupling data, but no chemical shift data. Later work by Lambert et al.¹³ and a detailed study by Canuel and St-Jacques¹⁴ again on deuterated-THP excluded the H3 and H4 chemical shifts. These analyses were in different solvents, CD₃OD/CHClF₂ and CS₂, with the former only giving $\Delta(\delta_{eq} - \delta_{ax})$ values. Eliel et al¹⁵ determined ²J_{HH} and ³J_{HH} and the R values obtained indicated that the THP ring was not flattened relative to cyclohexane. They also gave data for monosubstituted THP's but as unassigned multiplets of overlapping groups of protons. A more

recent study by Chu and True¹⁶ at 500 MHz assigned the room temperature ¹H chemical shifts as H3 1.657δ and H4 1.549δ despite an apparent peak intensity of 1:2. Thus even the combined published data does not provide a full assignment.

We give here the complete assignment of the proton spectrum of THP (**1**) at room temp. and below coalescence, the 3-methyl (**2**), 2-(hydroxymethyl) (**3**), 2-methoxy (**4**), 4-hydroxy (**5**), 4-(4'-hydroxybenzyl) (**6**) THP's and THP-4-carboxylic acid (**7**), 2-methyl-1,3-dioxolane (**8**) and two cyclic ethers of known conformation, 7-oxa-bicyclo[2.2.1] heptane (7-oxanorbornane) (**9**) and 1,8-cineole (**10**).



(9)



(10)

These plus previous literature results provides sufficient data for an analysis of oxygen SCS based on a previous theoretical model of proton chemical shifts¹. This model has been applied successfully to a variety of saturated hydrocarbons^{1,17} and their halogen¹⁸ and carbonyl¹⁹ derivatives. We shall show that this model provides a quantitative treatment for ether oxygen SCS and that these are due to electric field and steric effects of which the steric effect is the major contributor.

THEORY

As the theory has been detailed previously^{1,17} only a brief summary of the latest version (CHARGE5) is given here. The theory distinguishes between substituent effects over one, two and three bonds which are attributed to the electronic effects of the substituents and longer range effects due to the electric fields, steric effects and anisotropy of the substituents. The CHARGE scheme calculates the effects of atoms on the partial atomic charge of the atom under consideration, based upon classical concepts of inductive and resonance contributions.

If we consider an atom I in a four atom fragment I-J-K-L the partial atomic charge on I is due to three effects. There is an α effect from atom J given by the difference in the electronegativity of atoms I and J. A β effect from atom K proportional to both the

electronegativity of atom K and the polarisability of atom I. And a γ effect from atom L given by the product of the atomic polarisabilities of atoms I and L. The important carbon γ effect (i.e. C.C.C.H) is parametrised separately and is given by a simple $\cos\theta$ dependance where θ is the C.C.C.H dihedral angle. There are also routines for the methyl γ effect and for the decrease in the γ effect of the electronegative oxygen and fluorine atoms for CX_2 and CX_3 groups.

The total charge is given by summing these effects and the partial atomic charges (q) converted to shift values using eqn. 1

$$\delta = 160.84q - 6.68 \quad (1)$$

The effects of more distant atoms on the proton chemical shifts are due to steric, anisotropic and electric field contributions. H..H steric interactions were found to be shielding and X..H (X = C, F, Cl, Br, I) interactions deshielding according to a simple r^{-6} dependance (eqn 2).

$$\delta_{steric} = a_s / r^6 \quad (2)$$

Furthermore any X..H steric contributions on a methylene or methyl proton resulted in a push-pull effect (shielding) on the other proton(s) on the attached carbon.

The effects of the electric field of the C-X bonds (X= H,F,Cl,Br,I,C=O) were calculated from eqn 3 where A_Z was determined as 3.67×10^{-12} esu (63 ppm au) and E_Z is the component of the electric field along the C-H bond.

$$\delta_{el} = A_Z E_Z \quad (3)$$

For anisotropic groups the magnetic anisotropy contribution was calculated from the McConnell eqn involving the anisotropies of the magnetic susceptibilities of the group. It was only necessary to invoke this term for the unsaturated carbonyl group. The other groups investigated so far (C-X, X= H,F,C,Cl,Br,I) did not require this term and we will assume that this term is not significant for ether oxygen SCS (see later).

These contributions were then added to the shifts of eqn. 1 to give the calculated shift of eqn 4.

$$\delta_{total} = \delta_{charge} + \delta_{steric} + \delta_{anisotropy} + \delta_{el} \quad (4)$$

To apply the above theory to the ethers considered here the only addition necessary is the calculation of the electric field of the ether oxygen. The electric field for a univalent atom (e.g. fluorine) is calculated as due to the charge on the fluorine atom and an equal and opposite charge on the attached carbon atom. The vector sum gives the total electric field at the proton concerned and

the component of the electric field along the C-H bond considered is E_z in eqn 3. This procedure is both simpler and more accurate than the alternative calculation using bond dipoles.

For the ether oxygen two possible models were attempted. In the first the electric fields due to the charge on the oxygen atom and of both the attached carbon atoms were calculated at the proton in question. The carbon atoms each having a positive charge equal to half the oxygen charge. An alternative procedure was to calculate the electric field due to the ether oxygen atom and a dummy atom placed midway between the attached carbon atoms with an equal and opposite charge. As the coefficient in eqn 3 is known the electric field effect is given immediately without any further parametrisation. In practise the latter model gave better agreement with the observed data and this model is given here.

The steric effect of the oxygen atom is not known and therefore a value of the coefficient a_s in eqn 2 for oxygen must be determined. This and the associated push-pull coefficient are the only additional parameters required in order apply the theory to the more distant protons in ethers. The vicinal (H.C.C.X) effects are treated separately in CHARGE and for ethers an explicit H.C.C.O term equal $0.20\cos\theta$ was required (see later).

Experimental

THP (**1**), 3-methyl THP (**2**), 2-(hydroxymethyl)-THP (**3**), 2-methoxy-THP (**4**), 4-hydroxy-THP (**5**), 2-methyl-1,3-dioxolane (**8**), bicyclo-[2.2.1]-heptane (**9**) and 1,8-cineole (**10**) were obtained from Aldrich. 4-(4'-hydroxy-benzyl)-THP (**6**) and THP-4-carboxylic acid (**7**) were kindly donated by Zeneca Pharmaceuticals. **3** was distilled prior to use. The solvents were obtained commercially, stored over molecular sieves and used without further purification.

^1H and ^{13}C NMR spectra were obtained on a Bruker AMX400 spectrometer operating at 400.14 MHz for proton and 100.63 MHz for carbon. Spectra for **2** were also recorded on a Bruker DRX 600 spectrometer. Spectra were recorded in 10 mg cm^{-3} solutions (^1H) and ca. 50 mg cm^{-3} (^{13}C) with a probe temperature of ca. 25°C in CDCl_3 and referenced to TMS unless otherwise stated. Typical ^1H conditions were 128 transients, spectral width 3300 Hz, 32k data points, giving an acquisition time of 5s and zero-filled to 128k to give a digital resolution of 0.025 Hz.

The 2D experiments were performed on the AMX400 and Bruker DRX-600 using the standard Bruker COSY-DQF and HXCO-BI pulse sequences²⁰. The geometries of the compounds investigated were obtained by geometry optimizations using the GAUSSIAN94 programme at the

RHF/6-31G* level²¹. Full details of these optimizations and geometries are given in Ref. 22. The GAUSSIAN94 calculations were performed on the University of Liverpool Central Computing facility, and the CHARGE5 computations were performed on a PC.

Assignments

Tetrahydropyran (1). The room temperature ¹H spectra of THP in 50:50 v/v CDCl₃:CFCl₃ consists of 3 multiplets at δ 3.632, 1.637 and 1.568 of integration 2:1:2, assigned as H2, H4 and H3 respectively. At -85°C all proton chemical shifts are resolved with no change observed to -95°C. The H2 protons were at low field, and H3 and H4 distinguished by integration. Assignment of axial or equatorial was based upon the splitting patterns of the protons.

3-Methyl-tetrahydropyran (2). The H2 and H6 protons are to low field and the methyl doublet at δ 1.69. The COSY plot shows a connection from the methyl peak to H3ax which is on the low field side of the multiplet at *ca.* δ 1.48 to 1.75. This was distinguishable at 600 MHz as an axial pattern, and correlates to the low field axial proton at δ 3.00 (H2ax), which in turn assigns H2eq at δ 3.80. The two remaining low field protons are thus H6ax δ 3.32 and H6eq δ 3.80. H6ax strongly correlates to the centre of the multiplet at δ 1.62, which at 600 MHz is an axial pattern, indicating H5ax. The lone axial proton at δ 1.11 shows large couplings to both H3ax and H5ax assigning this as H4ax. H4ax correlates strongly to an equatorial proton at δ 1.81 which is assigned as H4eq. The high field side of the multiplet at δ 1.56 is thus H5eq. These assignments were confirmed by a HMQC experiment with the ¹³C assignments of Eliel et al.¹⁵, and are given in Table 1.

2-(Hydroxymethyl)-tetrahydropyran. (3) The low field equatorial pattern at δ 4.01 is assigned as H6eq. The remaining low field protons are in an overlapping multiplet of two axial protons and an AB type pattern. From the COSY correlations H6eq defines H6ax at δ 3.46, leaving H2ax at δ 3.44, and the AB pattern as non-degenerate CH₂ protons at *ca.* δ 3.53. H2ax correlates strongly to an axial pattern at δ 1.32 assigning this as H3ax. The lone equatorial pattern at δ 1.86 correlates to H3ax and H6eq ('W' coupling), but not H6ax, and is thus H4eq. The remaining protons (H3eq/H4ax/H5ax/H5eq) form a strongly overlapping multiplet at δ 1.50 to 1.57. Correlations to H3ax and H4eq show H3eq δ 1.53 and H4ax δ 1.51 to be at the high field end of this multiplet. H5ax and H5eq form a complex second order pattern centered at δ 1.56. These assignments were

confirmed by a HET-CORR experiment with the ^{13}C assignments of C2 δ 78.40, C3 δ 27.46, C4 δ 22.98, C5 δ 26.01, C6 δ 68.32 and CH_2OH δ 66.22. The hydroxyl proton position varied on successive experiments between δ 2.1 to 2.3, but was distinctive as a doublet of doublets.

2-Methoxy-tetrahydropyran.(4) The H2 and H6 protons are immediately assigned and the remaining protons assigned by the COSY connections of H2e/H3e and H4ax/H3e with no correlations of H4ax to H2,6. H3e and H4a can be clearly distinguished at 400MHz but the remaining protons are closely coupled and their chemical shifts are only accurate to 0.01ppm.

4-Substituted tetrahydropyrans. The room temperature spectra of **5**, **6** and **7** can be immediately assigned and the spectral data is given in Table 1. For **6** the aromatic spectrum was analyzed using the LAOCOON software²³ on the hydroxyl decoupled spectrum to give a rms of 0.090 Hz (and probable errors of 0.024 to 0.042 Hz), and the chemical shifts and couplings in Table 1.

2-methyl-1,3-dioxolane.(8) The A_2B_2 spectrum of the ring protons has been analysed previously at 60 MHz²⁴ but at this frequency only 16 of the 24 theoretically allowed transitions were observed and it was necessary to assume $J_{AA'} = J_{BB'}$. At 400MHz. all the allowed transitions were resolved and thus all the couplings could be obtained by iterative analysis. The results are given in table 1, the rms error (LAOCOON) was 0.009Hz with probable errors of 0.003Hz (couplings) and 0.009Hz (chemical shifts). Note that the assignment of the H4,5 protons (syn and anti w.r.t. the methyl group) cannot be obtained from the analysis but was made from the calculated shifts (see later).

7-Oxa-bicyclo[2.2.1]heptane.(9) The CH_2 protons occur at 1.69 and 1.15 δ , and the CH proton at 4.56 δ . The low field methylene proton was assigned as exo since it showed the largest coupling to the methine proton (shown by decoupling the CH proton).

1,8-Cineole.(10) The methyl resonances at δ 1.24 (C_{Me} δ 28.8) and δ 1.05 (C_{Me} δ 27.5) of integration 2:1 are assigned to 10/11-Me and 9-Me respectively. The multiplet at δ 1.41 consists of sets of overlapping triplets or quartets ($J \sim 1.2$ Hz) and with an integration of one proton is H4 (C_4 δ 32.8). From the COSY plot H4 couples to the multiplet at δ 1.5 (4 protons) and δ 2.0 (2 protons), but not at δ 1.7 (2 protons) which is thus a C2 proton. From the HET-CORR the C_3 (δ 22.7) has protons at δ 1.5 and 2.0, and C_2 (δ 31.4) at δ 1.5 and 1.7. The assignment of the protons for C_3 can be made on the basis of the coupling from H4 to δ 2.0 (3x) \gg δ 1.5 (3n). This assignment is consistent with that in bornanes with the exo proton to low field of the endo

protons¹. The multiplet pattern at δ 1.66 is similar to 3x and could tentatively be assigned as 2x, although the pattern at δ 1.5 is not obviously endo or exo. Consideration of chemical shifts might suggest $\delta(\text{exo}) > \delta(\text{endo})$ yet in 1,4-cineole²⁵ the opposite was found from lanthanide induced shift (LIS) analysis.

The assignments of **9** and **10** were confirmed by their LIS using Yb(FOD)₃. For **9** the assignment of the exo proton to low field was confirmed by the greater LIS than the endo, since the exo proton is closer to the lanthanide at the oxygen binding site. For **10** the assignments of 2x with the largest methine proton LIS shift, and 3x with a LIS greater than 3n, are confirmed.

Results

The low temperature chemical shift differences ($\delta_{\text{eq}} - \delta_{\text{ax}}$) of **1** derived from the data in Table 1 are in good agreement with the literature values. E.g. H2 0.558 cf. 0.55¹¹, 0.50¹³, 0.527¹⁴, 0.563¹⁶; H3 -0.116, cf. -0.074¹⁴ and H4 0.367 cf. 0.32¹³. $\delta_{\text{H3ax}} > \delta_{\text{H3eq}}$ in contrast to H2 and H4 (Table 1) and the general cyclohexane rule of $\delta_{\text{eq}} > \delta_{\text{ax}}$. This may explain why the average δ_{H4} is greater than δ_{H3} at room temperature against the assignment of Chu and True¹⁶.

For **2** ΔG° (eq-ax) is 1.44 kcal mol⁻¹¹⁵, thus ca 6% of the axial form is present at room temperature. This is sufficiently small that the resultant oxygen ring substitution shift should be relatively unaffected by the minor conformer. Note that $\delta_{\text{H5ax}} > \delta_{\text{H5eq}}$ confirming an orientational oxygen γ effect.

It is of interest to consider the oxygen SCS in these systems in more detail and Figure 1 shows the oxygen ring substitution effects of **1** relative to cyclohexane, **2** relative to methylcyclohexane^{1,17} and also the SCS of the 2-methoxy group in **4**. In THP the β effect of the oxygen atom on H2ax and H2eq is +2.1 to +2.3 ppm and non-orientational. The γ effect is strongly deshielding on H3ax (+0.4 to +0.5 ppm) but shielding on H3eq (ca. -0.1 ppm). This orientational effect is also seen in 1,3-dioxanes where the ring substitution shifts of 4-methyl-1,3-dioxane (table 2) vs. methylcyclohexane¹ on H5ax is +0.83 ppm and for H5eq -0.28 ppm. Here the effect of two oxygen atoms would appear roughly additive. Finally, the long range oxygen effect on H4 is deshielding at +0.15 to +0.3 ppm, with the greatest effect on the axial proton.

The oxygen ring substitution effect of 7-oxa-norbornane versus norbornane¹⁷ may be compared with the results for THP (**1**). For the methine proton the oxygen shift of +2.37 ppm is

in accord with the β THP protons. The oxygen SCS of + 0.22 ppm (Hexo) and -0.01 ppm (Hendo) again show the orientational oxygen SCSs, although less pronounced than for THP.


Table 1. ¹H Chemical Shifts (δ) and H-H Couplings (Hz) in THP (**1**), 3-methyl-THP (**2**), 2-(hydroxymethyl)-THP (**3**), 2-methoxy THP (**4**), 4-hydroxy-THP (**5**), 4-(4'-hydroxybenzyl)-THP (**6**), THP-4 carboxylic acid (**7**) and 2-methyl-1,3-dioxolane (**8**).

Proton	1 ^a	5 ^a	2 ^b	3 ^b	4 ^b	5 ^b	6 ^b	7 ^b	8 ^b	
		eq-OH	ax-OH							
2ax	3.439	3.407	3.836	2.971	3.437	-	3.444	3.340	3.457	4.987
2eq	3.997	4.029	3.75 ^c	3.801	-	4.517	3.962	3.954	3.990	-
3ax	1.682	1.579	d	1.686	1.323	1.60	1.566	1.315	1.803	-
3eq	1.566	1.915	d	-	1.53 ^e	1.732	1.904	1.551	1.885	-
4ax	1.493	3.751	-	1.114	1.51 ^e	1.816	3.860	1.692	2.593	3.857 (syn)
4eq	1.860	-	4.229	1.810	1.856	1.55	-	-	-	3.993 (anti)
5ax				1.618	1.56 ^e	1.60				3.857 (syn)
5eq				1.563	1.56 ^e	1.55				3.993 (anti)
6ax				3.316	3.463	3.856				
6eq				3.860	4.012	3.522				
CH ₃				0.805		3.405				1.384
CH ₂					3.569		2.470			
					3.507 ^f					
Ph							6.750			
							7.006			
H-H Couplings										
		5		6 ^h			7		8	
² J _{2ax2eq}		-11.29		-11.81			-11.58		² J _{HH}	-7.74
² J _{3ax3eq}		-13.21		-13.39			-13.60		³ J _{cis(AA')}	7.04
³ J _{2ax3ax}		9.85		11.87			10.86		³ J _{trans(AB')}	6.12
³ J _{2ax3eq}		2.68		2.15			2.69		³ J _{cis(BB')}	7.46
³ J _{2eq3ax}		4.01		4.53			4.26		³ J _(Me,H)	4.83
³ J _{2eq3eq}		4.45		g			3.00			
³ J _{3ax4ax}		9.03		11.48			10.82			
³ J _{3eq4ax}		4.43		3.75			4.32			

a) in CDCl₃:CFCl₃ at -85°C. b) in CDCl₃ at RT. c) obscured by H4ax of eq-OH. d) obscured by respective eq-OH protons. e) cf. ¹³C-¹H correlations. f) non-degenerate CH₂ group. g) long range 'W' couplings complicating 2eq and 3eq patterns.

h) Compound **6**, benzyl group δ (CH₂) 2.470, Ph 2,6H 7.006; 3,5H 6.750, couplings J₂₃ 8.14, J₂₄ 2.65, J_{3,5} 2.47, J₂₅ 0.45, J(CH₂,Ho) 0.53, J(CH₂,H4ax) 7.19.

Figure 1: a) The oxygen ring SCS for **1** and **2**, and b) the SCS of the 2-methoxy group in **4**.



Even more remarkable is the SCS of the 2-methoxy group in 2-methoxy-THP (figure 1b). The 2-methoxy group is predominantly axial (ΔG° (eq-ax) is ca $-0.8 \text{ kcal mol}^{-1}$)¹⁰ as shown in Figure 1. The values of the methoxy SCS are confirmed by the similar data for the trans-2-methoxy-4-methyl THP (Table 2) in which due to both the substituent groups being in their favourable conformation the molecule is entirely in the conformation with eq-methyl and axial methoxy. The corresponding values for the 2-methoxy SCS at the 6ax and 6eq protons are +0.38 and -0.34 ppm.¹⁰

This data together with the other data in Table 2 provides a rigorous test of the application of the theoretical model to these compounds. The six-membered ring compounds in Table 2 all exist predominantly in one chair conformation but the five membered rings of THF and 1,3 dioxolane exhibit pseudorotation. For THF both the O-envelope (C_2) and half-chair (C_s)

conformations were considered. (At the 6-31G* level of theory the Cs conformer was favoured by 0.39 kcal mol⁻¹). The ab initio calculations for dioxolane iterated preferentially to the Cs conformation which was used in the calculations. However the calculations for the 2-methyl-1,3-dioxolane gave an envelope conformation with C₂ out of the plane of the other ring atoms. The calculated shifts for the 2-methoxy-THP derivatives are given for the axial conformation of the methoxy group. In this conformation there are only two populated conformations of the methoxy methyl group as the conformer with the methyl over the THP ring (i.e. with the H.C₂.O.Me dihedral ca. 180°) is very unfavourable. The calculations for the two gauche conformers (H.C₂.O.Me dihedral) are given in Table 2, although the rotamer with ∠O.C.O.Me of -64° is preferred over ∠O.C.O.Me of 145° at RHF/6-31G* calculations by 4.0 kcal mol⁻¹.

As previously mentioned the electric field effect is given directly by equation 3 thus only the steric coefficient a_s (equation 2) and the push-pull coefficient need to be determined. The data in Figure 1 suggest that the push-pull coefficient for the methoxy group should be ca. -1.0 and this was confirmed in the calculations together with the appropriate value of a_s for oxygen as 100.0.

The observed vs. calculated shifts for a range of ethers is given in Table 2 and it is of some interest to consider these results. The general agreement of the observed vs. calculated shifts is very good and the great majority of the observed shifts are reproduced to better than 0.1 ppm. The agreement is particularly striking for the chair conformations of THP and 1,3-dioxane with an overall rms error (observed vs. calculated shifts) of < 0.05 ppm. In both cases the calculations reproduce all the oxygen SCS very well and in particular the low field shift of the axial proton w.r.t. the corresponding equatorial proton in 3a vs. 3e (THP) and 5a vs. 5e (3-methyl THP and 1,3-dioxanes) is strikingly reproduced. The only significant discrepancy for these compounds is H4ax for which the calculated shift is ca 0.2 ppm too low.

The agreement between the observed and calculated shifts though reasonable is not as good in the case of the boat structures of 7-oxanorbornane and 1,8-cineole. This may well be due to the fact that the proton shifts in the parent hydrocarbons of norbornane and bicyclo-2,2,2-octane are not as well reproduced as those of cyclohexanes in this theoretical model and these discrepancies may well carry over to the oxygen analogues. E.g. the observed shift of the bridgehead proton in norbornane is to low field of the calculated value (2.19 vs. 1.95 δ)¹ and the difference is a similar to the observed-calculated shift of H1 in 7-oxanorbornane (Table 2).

However even in these cases the model does provide a basis for estimating the proton shifts in these compounds and indeed the correct order of the proton shifts is given in every case.

The good agreement between the observed and calculated shifts over this diverse range of compounds is strong support for our original assumption to neglect any C-O bond anisotropy contributions and this is also in accord with Zurcher's earlier conclusions^{3a}.

The theoretical interpretation of these SCS's on the basis of the present theory is of some interest. The oxygen γ effect is deshielding for a gauche O.C.C.H orientation and shielding for a trans orientation, in contrast to the carbon γ effect which is shielding for a gauche C.C.C.H orientation and deshielding for the trans orientation. Thus the large differences in the oxygen γ effects shown in H3ax vs. H3eq in THP etc. are due to the replacement of the carbon at C1 by the oxygen atom. The consequent replacement of the carbon γ effect by the oxygen γ effect gives rise to the observed oxygen SCS.

The oxygen SCS at the more distant H4 proton (and H4 and H6 in 1,3-dioxane) are also due to the replacement of the CH₂ group interactions by the oxygen atom. In this case the interactions concerned are the steric and electric field terms and these have much less orientation dependence in these systems. Both terms are quite small at these protons, e.g. for H4 of THP the steric interactions with the oxygen atom are 0.03 and 0.08 ppm respectively for H4e and H4a and the corresponding electric field contributions are 0.13 and 0.12 ppm. The alternative model in which positive charges were placed on the carbon atoms attached to the oxygen atom gave generally smaller effects and poorer agreement with the observed shifts.

In contrast the steric term and the associated push-pull contribution dominate the SCS of the axial 2-methoxy group. E.g. for H4a the calculated contributions are 0.32ppm (steric) and 0.17ppm (electric field) and these well reproduce the observed shifts. There are similar calculated values for H6a but here the calculated shift is somewhat larger than the observed shift.

Table 2. Observed vs. calculated proton chemical shifts (δ) of acyclic and cyclic ethers.

Molecule		Experimental ^a	CHARGE5	
Dimethyl ether	Me	3.24 ^b	3.31	
Methyl- t-butyl ether	Me	3.22 ^c	3.43	
	t-Bu	1.19	1.27	
Tetrahydrofuran	α -CH ₂	3.83 ^c	3.80(C ₂), 3.84(Cs)	
	β -CH ₂	1.85	1.58(C ₂), 1.54(Cs)	
Tetrahydropyran	2e	4.00	3.99	
	2a	3.44	3.44	
	3e	1.57	1.54	
	3a	1.68	1.65	
	4e	1.86	1.80	
	4a	1.49	1.25	
3-Methyl-THP	2e	3.80	3.92	
	2a	2.97	3.10	
	3e (Me)	0.81	0.86	
	3a (CH)	1.69	1.85	
	4e	1.81	1.77	
	4a	1.11	0.87	
	5e	1.56	1.56	
	5a	1.62	1.64	
	6e	3.86	3.99	
	6a	3.32	3.47	
2-Methoxy-THP	2e (CH)	4.52	4.79 ^h	4.68 ⁱ
	2a (OMe)	3.41	3.37	3.32
	3e	1.73	1.77	1.72
	3a	1.60	1.53	1.41
	4e	1.55	1.53	1.53
	4a	1.82	1.74	1.75
	5e	1.55	1.57	1.57
	5a	1.60	1.64	1.62
	6e	3.52	3.64	3.61
	6a	3.86	4.19	4.21
Trans- 2-methoxy-4-methyl-THP				

	2e (CH)		4.64 ^d	4.76 ^h	4.70 ⁱ
	2a (OMe)		3.30	3.38	3.33
	3e		1.67	1.73	1.69
	3a		1.24	1.17	1.05
	4e (Me)		0.87	0.87	0.87
	4a (CH)		1.88	1.95	1.97
	5e		1.52	1.54	1.53
	5a		1.23	1.28	1.26
	6e		3.53	3.66	3.63
	6a		3.70	4.18	4.20
1,4-Dioxane	CH ₂		3.70 ^c	3.85	
1,3,5-Trioxane	CH ₂	5.00 ^b	5.15 ^c	4.93	
2,4,6-Trimethyl-1,3,5-trioxane	CH		5.05 ^e	4.59	
	Me		1.40	1.37	
1,3-Dioxane	2-H		4.84 ^c	4.85	
	4-H		3.90	3.73	
	5-H		1.78	1.75	
2-Methyl-1,3-dioxane	2e (Me)		1.15 ^f	1.29	
	2a (CH)		4.50	4.59	
	4e		3.95	4.17	
	4a		3.61	3.52	
	5e		1.23	1.44	
	5a		1.95	2.12	
4-Methyl-1,3-dioxane	2e	5.05	4.89 ^g	5.19	
	2a	4.71	4.55	4.53	
	4e (Me)	1.24	1.17	1.27	
	4a (CH)	3.73	3.61	3.59	
	5e	1.48	1.38	1.42	
	5a	1.76	1.69	1.67	
	6e	4.09	4.02	4.15	
	6a	3.71	3.58	3.53	
2-methyl-1,3-dioxolane	2(Me)		1.38	1.30	
	2(CH)		4.99	4.75	
	4,5 (anti)		3.99	3.99	
	4,5 (syn)		3.86	3.79	
7-Oxa-norbornane	H1 (CH)		4.56	4.17	

	Hex	1.69	1.86
	Hen	1.44	1.30
1,8-Cineole	2ex	1.66	2.00
	2en	1.50	1.28
	3ex	2.02	1.90
	3en	1.50	1.32
	H4 (CH)	1.41	1.38
	9-Me	1.05	1.18
	10/11-Me	1.24	1.19
1,3-Dioxolane	α -CH ₂	4.90 ^c	5.02
	β -CH ₂	3.88	3.89

a) This work unless stated. b) Ref. 26. c) Ref. 27. d) Ref 8. e) Ref. 28. f) Ref 4. g) Ref 5. h) O.C.O.Me dihedral -64° . i) O.C.O.Me dihedral 145° .


As a further check on the accuracy of these calculations Table 3 gives the calculated δ values for 4-oxa-5- α -androstande and also the comparable values for the parent 5- α -androstande from which the calculated SCS can be obtained to compare with the observed SCS from Ref. 4. The agreement is very good and again the vicinal equatorial proton in the tetrahydropyran ring (2α in Table 3) shows an upfield SCS. The only disagreement between the observed and calculated SCS occurs for the 1α and 6 protons. The 1α proton is exactly analogous to H4a of THP and the 6 protons in 5- α -androstande are part of a complex unresolved multiplet with the 4 protons and thus the observed SCS will not be known very accurately¹.

Table 3. Calculated chemical shifts for 5 α -androstande and 4-oxa-5 α -androstande and calculated vs. observed SCS.^a

Proton	Calculated		SCS	
	Androstane ^b	4-Oxa-	Calc.	Obs. ^c
1 α	1.00	1.09	0.09	0.23
1 β	1.53	1.69	0.16	0.08
2 α	1.57	1.45	-0.12	-0.10
2 β	1.44	1.90	0.46	0.44
3 α	1.19	3.42	2.23	2.21
3 β	1.74	4.08	2.34	2.32
5 (CH)	1.15	3.27	2.12	1.88
6 α	1.35	1.95	0.60	0.37
6 β	1.40	1.92	0.52	0.21
7 α	0.97	0.95	-0.02	0.06
7 β	1.94	1.96	0.02	0.05
11 α	1.51	1.54	0.03	-0.03
11 β	1.34	1.47	0.13	0.05
12 α	1.05	1.06	0.01	-0.01
12 β	1.60	1.63	0.03	0.00
14 (CH)	0.83	0.83	0.00	-0.03
19-Me	0.80	0.94	0.14	0.15

a) protons with SCS < 0.01 ppm excluded. b) cf. Ref. 1. c) observed SCS cf. 4-oxa-androstan-17-one vs. androstan-17-one, Ref. 4.

Figure 2. Nomenclature used for 4-oxa-5 α -androstande.



Conformational Analysis

4-Hydroxy-THP (5). The value of ${}^3J_{2ax3ax}$ of 9.85 Hz in **5** (Table 1) suggested a significant percentage of the axial OH conformer at room temperature (vs. 11.87 Hz for **6**) and this was confirmed by a low temperature experiment. Integration of H4ax in $CDCl_3:CFCl_3$ at $-85^\circ C$ gave 8.5% of the axial conformer which gives $\Delta G^\circ(eq-ax)$ equal $1.02 \text{ kcal mol}^{-1}$. This is almost identical to the ΔG° value of the hydroxyl group in cyclohexane⁸ ($1.04 \text{ kcal mol}^{-1}$) showing clearly that there is no OH...O interaction in this molecule.

2-(Hydroxymethyl)-THP (3). Compound **3** is a model for the $CH_2OH\cdots O$ interactions in sugars such as D-glucose and derivatives. The non-degeneracy of the methylene protons of the CH_2OH fragment in **3** suggests an interaction between the hydroxyl group and the THP ring oxygen, causing energy differences between the three staggered rotamers (gg, gt and tg using the nomenclature of Ref. 30, see Figure 3). The chemical shifts and couplings of this fragment in both a non-polar solvent ($CDCl_3$) and a polar solvent (d_6 -acetone) obtained from the LAOCOON analysis are given in Table 4. From the two couplings (${}^3J_{A,2ax}$ and ${}^3J_{B,2ax}$) it is possible to calculate the relative populations of the three rotamers from eqns 4 and 5 if the couplings in the individual conformers are known.

$$J_{obs.} = \sum_{i=1,3} n_i J_i \quad (4)$$

$$\sum_{i=1,3} n_i = 1 \quad (5)$$

The conformer couplings derived from PCModel calculations²⁹ on the HF/6-31G* geometries²¹ (Scheme 1) are compared to the values using the equations of Ref. 31 (Scheme 2) in Table 5, along with their resultant populations. The consistency between the two schemes is encouraging, although Scheme 2 appears to give more realistic values of the tg conformer. In chloroform the gt conformer is favoured over the gg form in about a 2:1 ratio, with almost none of the tg form present. In both the gg and gt conformers the possibility of intra-molecular hydrogen bonding between the hydroxyl proton and the ring oxygen exists, as shown in Figure 4. From ab initio calculations at the MP2/6-31G* level the $\underline{OH}\cdots O$ distance for the gg form is 2.728 Å and the gt conformer at 2.714 Å, suggesting for the latter isomer this interaction may be larger. This could account for some of the preference for the gt form.

Table 4: Temperature and solvent effects on the HH couplings (Hz) and ^1H chemical shifts (δ) of the CH.CH₂OH fragment in **3**.

Solvent	Temp/K	$\delta(2a)$	$\delta(A)$	$\delta(B)$	$\delta(\text{OH})$	J_{AB}	$J_{A,2aX}$	$J_{B,2aX}$	$J_{A,\text{OH}}$	$J_{B,\text{OH}}$	rms
CDCl ₃	298	3.437	3.569	3.507	2.181	-11.45	3.11	7.38	8.08	4.19	0.053
	233	3.485	3.606	3.543	2.494	-11.62	2.75	7.96	8.61	3.72	0.072
d ₆ -Acetone	288	3.301	3.379	3.441	3.524	-11.26	4.30	6.20	7.02	5.20	0.064
	223	3.293	3.369	3.436	4.270	-11.39	4.26	6.58	7.31	5.25	0.035

Table 5: Conformer couplings (Hz) and populations for the CH.CH₂OH fragment of **3**.

Conformer		gg	gt	tg
Coupling (Hz)				
Scheme 1	J(A,2aX)	3.67	2.97	10.68
	J(B,2aX)	0.60	10.73	4.70
Scheme 2	J(A,2aX)	2.50	3.10	10.70
	J(B,2aX)	0.60	10.70	5.20
Population (%)				
CDCl ₃	Scheme 1	34	74	-1
	Scheme 2	31	66	3
d ₆ -Acetone	Scheme 1	36	50	14
	Scheme 2	34	47	19

If intra-molecular hydrogen bonding is important in the determining the rotamers of **3**

then in a solvent which can hydrogen bond to the hydroxyl group (but not the ring oxygen) the tg form should be stabilised. Indeed, in acetone (see Table 5) the tg conformer is present at 15-20%, whilst the gt form decreases. The gg form is approximately the same in both chloroform and acetone solutions.

In the assignment given in Table 4 we note that the chemical shifts of A and B have reversed order from chloroform to acetone. In the alternative assignment with $\delta(A) > \delta(B)$ and $J_{A,2a} = 6.2$ Hz and $J_{B,2a} = 4.3$ Hz in acetone at 288 K, then using Scheme 1 the calculated populations are gg = 41%, gt = 21% and tg = 38%. The same trends are seen although the increase in the tg and the decrease in the gt conformers is more dramatic.

It is interesting to compare the result in **3** of $gt > gg > tg$ with D-glucosides where in general $gg \sim gt \gg tg$. In both cases the tg form is small (usually <5% for sugars in polar solvents) and this has been attributed to the destabilization of the tg conformer due to syn-diaxial (C6-OR...C4-OR) oxygen-oxygen interactions³². However in **3** there is no C4-oxygen and thus the stabilisation of the gt and gg conformers by intra-molecular hydrogen bonding is significant, and should also be considered relevant to sugars.

Acknowledgements

We thank ZENECA Pharmaceuticals Ltd for a fully funded research studentship (M.A.W.) and for samples of **5** and **6**. We are pleased to acknowledge the assistance of Dr. P. D. Mallinson and the University of Liverpool central computing facility for the operation of GAUSSIAN94.

Figure 3: The three staggered rotamers about the C(2)-C(MeOH) bond in **3**.





Figure 4: Possible intra-molecular hydrogen bonding (marked with a dashed line) in **3**.



References

1. Part 11, R.J. Abraham, L. Griffiths and M.A. Warne, *Magn. Reson. Chem.*, in press.
 2. D.M. Grant and R.K. Harris (Ed.), *Encyclopedia of NMR*, J. Wiley, N.Y. 1996.
 3. a) R.F. Zurcher, *Prog. Nucl. Mag. Res. Spectrosc.*, 1967, **2**, 205.
b). H.-J. Schneider, U. Buchheit, N. Becker, G. Schmidt and U. Siehl, *J. Am. Chem. Soc.*, 1985, **107**, 7027.
c). L.D. Hall, *Tetrahedron Lett.*, 1964, **25**, 1457.
 4. Y. Yang, T. Haino, S. Usui and Y. Fukazawa, *Tetrahedron*, 1996, **32**, 2325.
 5. C.P. Rader, *J. Am. Chem. Soc.*, 1969, **91**, 3248.
 6. H.R. Buys and E.L. Eliel, *Tetrahedron Lett.*, 1970, 2779.
 7. K. Pihlaja and P. Ayras, *Acta Chem. Scand.*, 1970, **24**, 531.
 8. E.L. Eliel, S.H. Wilen and L.N. Mander, *Stereochemistry of Organic Compounds*, J. Wiley, N.Y. 1994.
 9. J. Delmau, J.C. Duplan and M. Davidson, *Tetrahedron*, 1967, **23**, 4371.
 10. H. Booth, K.A. Khedhair and S.A. Readshaw, *Tetrahedron*, 1987, **43**, 4699.
 11. G. Gatti, A.L. Segre and C. Morandi, *J. Chem. Soc. B*, 1967, 1203.
 12. J.B. Lambert, R.G. Keske and D.K. Weary, *J. Am. Chem. Soc.*, 1967, **89**, 5921.
 13. J.B. Lambert, C.E. Mixan and D.H. Johnson, *J. Am. Chem. Soc.*, 1973, **95**, 4634.
 14. L. Canuel and M. St.-Jacques, *J. Org. Chem.*, 1976, **41**, 1380.
 15. E.L. Eliel, K.D. Hargrove, K.M. Pietrusiewicz and M. Manoharan, *J. Am. Chem. Soc.*, 1982, **104**, 3635.
 16. P.-S. Chu and N.S. True, *J. Phys. Chem.*, 1985, **89**, 2625.
 17. R.J. Abraham, L. Griffiths and M.A. Warne, *J. Chem. Soc. Perkin Trans. 2*, 1997, 31.
 18. (a) R.J. Abraham, L. Griffiths and M.A. Warne, *J. Chem. Soc. Perkin Trans. 2*, 1997, 203.
(b) *ibid* 1997, 881 (c) *ibid* 1997, 2151.
 19. R.J. Abraham and N.J. Ainger, ms in preparation.
 20. Bruker UXNMR version 010892, Bruker Meßtechnik GmbH, NMR Software Development, Silbersteifen, D-7512 Rheinstetten.
 21. Gaussian 94, Gaussian Inc., Pittsburgh PA, 1994.
- M.J. Frisch, G.W. Trucks, M. Head-Gordon, P.M.W. Gill, M.W. Wong, J.B. Foresman, B.G. Johnson, H.B. Schlegel, M.A. Robb, E.S. Replogle, R. Gomperts, J.L. Andres, K.

Raghavachari, J.S.Binkley, C.Gonzalez, R.L.Martin, D.J.Fox, D.J.Defrees, J.Baker, J.J.P.Stewart and J.A.Pople.

22. M.A.Warne, Ph.D. thesis, University of Liverpool, 1997.

23. S.M.Castello and A.A.Bothner-By, *J. Chem. Phys.*, 1967,**41**, 3863.

24. (a) R.J.Abraham, *J.Chem.Soc.* 1965, 256.

(b) R.J.Abraham, *Analysis of High Resolution NMR Spectra*, Elsevier Pub. Cor. , 1971.

25. W-G Liu, J.J.Steffens, A.Goswami and J.P.N.Rosazza, *J. Org. Chem.*, 1993, **58**, 6329.

26. J.W.Emsley, J.Feeney and L.H.Sutcliffe, *High Resolution NMR Spectroscopy*, Vol. 2, Appendix B, Pergamon Press, Oxford , 1966.

27. C.J.Puchert and J.Behnke, *Aldrich Library of ¹³C and ¹H FT NMR Spectra*, Aldrich Chemical Company Inc. 1993.

28. N.F.Bhacca, *Varian NMR Spectra Catalogue*, Varian Assoc. 1962.

29. PCModel version 5, Serena Software Ltd., PO BOX 3076, Bloomington, Indiana, 1994.

30. R.J.Abraham, E.J.Chambers and W.A.Thomas, *Magn. Reson. Chem.*, 1992, **30**, S60.

31. R.J.Abraham, J.Fisher and P.Loftus, *Introduction to NMR Spectroscopy*, pp.42-6, J. Wiley 1988.

32. Y.Nishida, H.Ohrui and H.Meguro, *Tetrahedron Lett.*, 1984, **25**, 1575.